

# The effect of doubled CO<sub>2</sub> and model basic state biases on the monsoon-ENSO system. Part A: Mean response and interannual variability

By A. G. TURNER\*, P. M. INNESS and J. M. SLINGO  
*Walker Institute, University of Reading, UK*

(submitted 10 August 2006, revised 21 January 2007)

## Summary

The impact of doubled CO<sub>2</sub> concentration on the Asian summer monsoon is studied using a coupled ocean-atmosphere model. Both the mean seasonal precipitation and interannual monsoon variability are found to increase in the future climate scenario presented. Systematic biases in current climate simulations of the coupled system prevent accurate representation of the monsoon-ENSO teleconnection, of prime importance for seasonal prediction and for determining monsoon interannual variability. By applying seasonally varying heat flux adjustments to the tropical Pacific and Indian Ocean surface in the future climate simulation, some assessment can be made of the impact of systematic model biases on future climate predictions. In simulations where the flux adjustments are implemented, the response to climate change is magnified, with the suggestion that systematic biases may be masking the true impact of increased greenhouse gas forcing. The teleconnection between ENSO and the Asian summer monsoon remains robust in the future climate, although the Indo-Pacific takes on more of a biennial character for long periods of the flux adjusted simulation. Assessing the teleconnection across interdecadal timescales shows wide variations in its amplitude, despite the absence of external forcing. This suggests that recent changes in the observed record cannot be distinguished from internal variations and as such are not necessarily related to climate change.

**This is a preprint of an article published in the Q. J. R. Meteorol. Soc. 133, 622, pgs 1143-1157.**

**<http://dx.doi.org/doi:10.1002/qj.82>**

**<http://www3.interscience.wiley.com/journal/114286409/abstract>**

**Copyright © Royal Meteorological Society 2007**

**Keywords:** India Predictability Interdecadal Variability Systematic Error El Niño

## 1. Introduction

Climate change will have important repercussions for the monsoon climates of Southeast Asia for a number of reasons. Not only could increased greenhouse gas forcing affect the mean monsoon, but further extremes of variability could have disastrous effects. More irregular or severe monsoon seasons could lead to water scarcity, food shortages and malnutrition during drought or an increase in water-borne disease after heavy rain, together with damage to local infrastructure.

Whilst making predictions of future monsoon behaviour is a key goal of climate change research, the findings are not always in agreement. May (2004) cites a number of different studies which give no overall indication of how the Asian summer monsoon will respond to increased greenhouse forcing. There are, however, a number of common themes. An increase in the land-sea meridional gradient of temperature in the Indian region may act to strengthen monsoon dynamics, as argued by Hu *et al.* (2000), or alternatively move the flow structure northward with no overall strengthening (Kitoh *et al.* 1997; Ashrit *et al.* 2003; May 2004). Most previous work sees an increase in seasonal rainfall over India, ranging from just the peninsula (Hu *et al.* 2000), to a varied response of increases in the northwest and far south (Ashrit *et al.* 2003), and an overall intensification of the mean pattern (Kitoh *et al.* 1997; May 2004). The warmer Indian Ocean and enhanced low-level moisture convergence have been found to play a major role, allowing increased seasonal precipitation even in regions of weaker flow (*e.g.*, Kitoh *et al.* 1997; Meehl *et al.* 2000; Ashrit *et al.* 2003).

In addition to the differential warming of land and sea, some studies such as Meehl and Washington (1996) notice a pattern of El Niño-like change in the equatorial Pacific (*i.e.*, the east warming more so than the west). Since El Niño is often associated with weak monsoons (*e.g.*, Webster *et al.* 1998), this pattern of change may push the Indian monsoon toward drought conditions,

\* Corresponding author: Walker Institute for Climate System Research, Department of Meteorology, University of Reading, PO Box 243, Earley Gate, Reading, RG6 6BB, UK. email: a.g.turner@rdg.ac.uk

via the atmospheric teleconnection. Indeed in a study of two coupled models under increasing CO<sub>2</sub> conditions, Meehl *et al.* (2000) noted that El Niño-like warming in one model led to a smaller increase in seasonal mean precipitation when compared to the other.

Aside from the mean response of the tropical Pacific, changes to ENSO variability would likely impact on the monsoon system. The understanding of future monsoon-ENSO behaviour can be helped by looking at current observations. Kumar *et al.* (1999) demonstrate that the inverse monsoon-ENSO relationship has broken down over recent decades, associated with a south-eastward shift in Walker circulation anomalies, reducing the anomalous subsidence over India and favouring normal monsoon conditions during El Niño. However, Torrence and Webster (1999) suggest that rather than a climatic shift, the weakened relationship is actually part of a series of variations on decadal timescales. They point to intervals of high (1875-1920, 1960-1990) and low (1920-1960) variance of the monsoon-ENSO system. During high periods, the system is highly coherent, lending itself to increased predictability. However, in experiments with an atmosphere-only GCM, Meehl and Arblaster (2003) demonstrate that even by fixing ENSO variability, monsoon variability can still increase under stronger greenhouse forcing due to changes in the basic state.

Although there are several large uncertainties contributing to the spread in future climate prediction (see, *e.g.*, Ashrit *et al.* 2003), this study particularly relates to biases in the mean state in GCMs and their effect on monsoon prediction. Just as Sperber and Palmer (1996) noted the link between correct simulation of the climatological basic state in the tropics and accurate seasonal prediction of precipitation variability (as confirmed in Turner *et al.* 2005, hereafter TIS05), Federov and Philander (2000), amongst others, realised that such errors could have an enormous effect on future predictions. The effect of these errors will be assessed here through analysis of two future-climate integrations of the Met Office Unified Model HadCM3, using both the standard version of the model and the equatorial Indo-Pacific flux corrected version (HadCM3IPFA) presented in TIS05. These experiments and data used are outlined in section 2. Section 3 considers the effect of climate change on the mean state, whilst section 4 addresses monsoon variability. These results are related to ENSO changes in section 5, whilst the predictability of the monsoon via its teleconnection to the Pacific Ocean is considered in section 6. Conclusions are drawn in section 7.

## 2. Model details and observed datasets

### (a) *The coupled model*

The GCM employed in this study is the fully coupled Unified Model, HadCM3, combining state-of-the-art atmosphere (Pope *et al.* 2000) and ocean (Gordon *et al.* 2000) components. HadCM3 was put to extensive use to provide climate change scenarios for the Intergovernmental Panel on Climate Change Third Assessment Report (IPCC 2001). Multi-century integrations can be performed with no significant climate drift, without the use of flux adjustments (Johns *et al.* 2003).

The atmospheric component (HadAM3) is solved on a regular grid of 3.75° longitude by 2.5° latitude. Whilst the default vertical resolution is 19 levels, 30 levels are used in this study, as in TIS05, halving the layer thickness in the mid-troposphere to 50mb. Inness *et al.* (2001) showed this configuration to give an improved representation of intraseasonal variability in tropical convection, such as the Madden Julian Oscillation (MJO), and a subsequent study (Spencer and Slingo 2003) found precipitation to respond more realistically to SST forcing, particularly El Niño. The oceanic component (HadOM3) has a uniform resolution of 1.25° by 1.25° with 20 levels in the vertical. The accurate simulation of the spatial structure and seasonality of the Asian summer monsoon (Martin *et al.* 2000) and of the monsoon trough (Johns *et al.* 2003), make HadCM3 well-suited to this study.

### (b) *Model integrations*

This study uses results from the HadCM3 L30 control run (pre-industrial CO<sub>2</sub>) presented in TIS05. The model was initialised from an L19 integration of HadCM3, and an initial period of 10 years was discarded to allow any spin-up to the increased vertical resolution. As noted in TIS05, the L30 version of the model is also stable and its climatology is similar to that of the standard L19 version.

The HadCM3 2 × CO<sub>2</sub> experiment was initialised from an existing 150-year 2 × CO<sub>2</sub> run, previous to which the carbon dioxide concentration had been ramped up at a rate of 1%.year<sup>-1</sup> for 70 years. Although unlikely to have achieved true equilibrium, the 150-year stabilization period prior to measurements being taken is in accordance with IPCC guidelines (see, *e.g.*, Merryfield 2006).

The first 10 years of the new experiment were discarded as spin-up to L30, and the integration was continued for a further 100 years. This experiment is designed to look at the effect of climate change on the monsoon climate, the ENSO system and the teleconnection between them. The non-transient nature of the forcing allows the monsoon-ENSO system to be studied in detail, including its interdecadal variability.

HadCM3, as with all coupled models, contains systematic biases under control conditions (Inness *et al.* 2003; TIS05), as well as in future greenhouse forcing scenarios. However, there is no way of ascertaining whether the adjustments to counteract those errors would be the same in a future climate simulation. Consequently the same seasonal cycle of flux adjustments (as used in TIS05 to correct biases in the mean state) are applied to a future climate integration, known hereafter as HadCM3IPFA  $2 \times \text{CO}_2$ . Heat fluxes only are fully applied in the equatorial band ( $\pm 5^\circ\text{N}$ ) of the Indian and Pacific Oceans, and are ramped down to zero at  $\pm 10^\circ\text{N}$ . Previous climate change experiments with older coupled GCMs kept the same seasonal cycle of anomalous fluxes as in their current climate (*e.g.*, Collins 2000a, using HadCM2). The objective of the HadCM3IPFA  $2 \times \text{CO}_2$  experiment is to explore the potential uncertainty caused by systematic biases rather than to necessarily give a better future prediction than HadCM3. The results of a 95-year integration of this model are presented.

The integrations will be referred to in the text as HadCM3 and HadCM3IPFA, both under  $1 \times \text{CO}_2$  and  $2 \times \text{CO}_2$  conditions. Neither integration takes account of sulphate aerosol forcing.

### (c) Observed datasets

Observational and reanalysis datasets have been used in this study as a means of comparison with the current climate. Monsoon variability has been assessed using the ECMWF reanalysis dataset (ERA-40, Uppala *et al.* 2005) for the dynamical fields, the CMAP dataset (Xie and Arkin 1997) and All-India Rainfall (AIR, Parthasarathy *et al.* 1994) for precipitation. Long-term El Niño behaviour has been assessed using HadISST data 1900-1999 (Rayner *et al.* 2003).

## 3. The influence of climate change on the GCM basic state

### (a) HadCM3

The effect of climate change (doubling  $\text{CO}_2$  concentration) on the mean summer climate of HadCM3 is shown in Fig. 1. Firstly, considering surface temperatures, (Figs. 1a,d) it is clear that warming has taken place across the whole of the Indo-Pacific sector. In the Pacific this is manifest as El Niño-like warming (*i.e.*,  $+1.5$  to  $2^\circ\text{C}$  in the east, compared to  $+1^\circ\text{C}$  in the west). This mean response amounts to a reduction in the zonal equatorial temperature gradient, and may lead to suppression of the monsoon following the arguments of Meehl *et al.* (2000). Such warming patterns were also noted by Timmermann *et al.* (1999) in their transient climate change integration, and by Ashrit *et al.* (2003) who found it to peak during boreal summer. However, in their review of the CMIP† integrations, Collins *et al.* (2005) found the most likely scenario was one of no El Niño- (or La Niña-) like climate change. Indeed, those models with the largest El Niño-like warming in future climate scenarios were generally those featuring poorly simulated present day ENSO variability. As an exception HadCM3 was found to exhibit realistic ENSO variability when compared to other coupled models (Latif *et al.* 2001). Such a warming pattern on the surface complements changes in equatorial subsurface waters, where the vertical temperature gradient is enhanced, amounting to an intensification of the thermocline, and deepening in the east (not shown). Collins (2000b) attributes the increased vertical temperature gradient at  $2 \times \text{CO}_2$  to the slow penetration of heat into the ocean. Once warmed, the upper levels become stable due to their buoyancy, inhibiting mixing and warming at depth.

Figure 1d shows land regions have warmed to a greater degree than the ocean. Given this is an equilibrium experiment, the explanation is not as simple as the much larger heat capacity of water. Sutton *et al.* (2007) have suggested that, assuming the anomalous surface climate forcing is evenly distributed over land and sea, the upward energy flux required to balance this will consist of a greater latent heat flux over the ocean. Thus temperatures over the land are more easily raised as there is less potential for increased evaporation, leading to a land-sea warming contrast. Relatively more warming occurs in the northern hemisphere due to the uneven distribution of the continents, as noted by, *e.g.*, Timmermann *et al.* (2004). This differential in surface warming has led to an increase in the land-sea meridional gradient of temperature in the Indian region, which may act to strengthen the monsoon

† Coupled Model Inter-comparison Project




Figure 1. HadCM3 2 × CO<sub>2</sub> summer (JJAS) climatologies of (a) surface temperature (K), (b) 850hPa winds (ms<sup>-1</sup>) and (c) precipitation (mm.day<sup>-1</sup>). Panels (d) to (f) show differences in these fields from HadCM3 1 × CO<sub>2</sub>. Unit wind vectors are 20ms<sup>-1</sup> (climate) and 5ms<sup>-1</sup> (anomaly). Speckling on the precipitation difference (f) indicates significance at the 95% level using a student t-test.

dynamics (Hu *et al.* 2000). However the lower tropospheric winds (850hPa) shown in Figs. 1b,e suggest that increased greenhouse forcing has affected the mean winds in only a few localities. The Pacific trade winds at about  $5^{\circ}\text{N}$  are approximately  $1\text{ms}^{-1}$  weaker in the region  $150^{\circ}\text{E} - 150^{\circ}\text{W}$ , consistent with the El Niño-like surface warming. Over the Indian Ocean, the flow at lower levels is more easterly just west of Sumatra, *i.e.* weaker across the south Bay of Bengal. It is difficult to relate this weakening to Pacific El Niño-like warming given that it is accompanied by stronger flow ( $+1\text{ms}^{-1}$ ) in the northern Arabian Sea and the north of India. This amounts to a slight northward shift of the monsoon flow structure, attributed to the increased meridional heating gradient by May (2004).

The anomalous easterlies out of Sumatra are part of a small anomalous anticyclone in the southern hemisphere, leading to local drying (of up to  $3\text{mm}\cdot\text{day}^{-1}$  in Fig. 1f). Significant increases (at the 95% level using a student t-test) in mean precipitation are noted in the main summer ITCZ and SPCZ bands across the Pacific Ocean. The northward displacement of the ITCZ is related to the preferential warming of the northern hemisphere, displacing the thermal equator in HadCM3 (Johns *et al.* 2003). Considering the Asian Summer Monsoon region, it is clear that the mean precipitation during the summer season has increased by between  $0.5\text{mm}\cdot\text{day}^{-1}$  in the northern states and at the tip of peninsula India, and  $2\text{mm}\cdot\text{day}^{-1}$  in the north-eastern Bay of Bengal, Bangladesh and Myanmar. The South China Sea also sees significant increases of this magnitude.

The response of HadCM3 to  $2 \times \text{CO}_2$  compares well with other studies. In agreement with work cited in the introduction (Kitoh *et al.* 1997; Meehl *et al.* 2000; Hu *et al.* 2000; Ashrit *et al.* 2003; May 2004), HadCM3 shows an overall increase in monsoon precipitation. However the spatial pattern of change is most similar to that found by Meehl and Arblaster (2003) using the Parallel Climate Model (PCM), although they found no drying south-west of Sumatra. Rather than the increased meridional temperature gradient, they attribute the increase in mean south Asian monsoon rainfall to the warmer Indian Ocean providing increased moisture, as in an earlier study by Meehl *et al.* (2000). Enhanced moisture convergence is also cited as the principal mechanism providing increased precipitation by Kitoh *et al.* (1997) and Ashrit *et al.* (2003). In this modelling framework, measurements of surface evaporation change between  $1 \times \text{CO}_2$  and  $2 \times \text{CO}_2$  are minimal (not shown), demonstrating that precipitation increases in HadCM3 are due to increased moisture convergence. The small response of the monsoon circulation to increased  $\text{CO}_2$  suggests non-dynamical causes for the increased moisture (Ashrit *et al.* 2003), and increased atmospheric water content over the northern Indian Ocean in HadCM3  $2 \times \text{CO}_2$  (not shown) confirms this.

Using SRES transient simulations in HadCM3, Johns *et al.* (2003) also noted significantly wetter conditions for South Asia during summer, although a weaker monsoon response was found when sulphate forcing was incorporated into the B2 scenario, consistent with the cooling effects of aerosols. The inclusion of aerosol forcing was also found to partially counter the effects of increased  $\text{CO}_2$  on monsoon precipitation in a study by Ashrit *et al.* (2003), highlighting the uncertainty involved in projections of future climate.

### (b) HadCM3IPFA

Figure 2 shows the mean summer climate of the HadCM3IPFA  $2 \times \text{CO}_2$  integration (a-c) and its differences with HadCM3IPFA  $1 \times \text{CO}_2$  (d-f). Looking at the summer (JJAS) surface temperature difference in Fig. 2d, it is clear that the flux corrected version of the model shows a similar response pattern to the standard model (Fig. 1d), with increased warming over land regions and a preferential response in the East Pacific, reducing the east-west SST gradient. This is accompanied by vertical intensification of the thermocline and deepening in the east, as in HadCM3 (not shown).

The surface wind response shown in Fig. 2e indicates a more marked pattern of change in the Indian monsoon region than in HadCM3 (Fig. 1e), with the increase in the low-level flow over the northern Arabian Sea being more pronounced. There is increased convergence at 850hPa in the warmer eastern equatorial Pacific, which aids the increased precipitation along the ITCZ noted in Fig. 2f. The magnitude of rainfall increase in the tropical convergence zone of the Pacific is much larger with flux adjustments, reaching almost  $5\text{mm}\cdot\text{day}^{-1}$  with climate change. The region of drying in HadCM3 west of Sumatra (Fig. 1f) is slightly larger and farther west, likely related to the stronger anticyclone there. In the Indian region, the same response pattern is represented in both versions of the model, although here it is stronger, especially in the north of India, the north Bay of Bengal, Bangladesh and south-central China.

The patterns of change are very similar in all three fields shown in Figs. 1 and 2, although the flux adjusted version seems to exhibit a stronger response to climate change. The only major difference




Figure 2. HadCM3IPFA  $2 \times \text{CO}_2$  summer (JJAS) climatologies of (a) surface temperature (K), (b) 850hPa winds ( $\text{ms}^{-1}$ ), and (c) precipitation ( $\text{mm}\cdot\text{day}^{-1}$ ). Panels (d) to (f) show differences in these fields from HadCM3IPFA under pre-industrial conditions. Unit wind vectors are  $20\text{ms}^{-1}$  (climate) and  $5\text{ms}^{-1}$  (anomaly). Speckling on the precipitation difference (f) indicates significance at the 95% level using a student t-test.




Figure 3. Response of the mean summer (JJAS) climate to flux adjustments at  $2 \times \text{CO}_2$  for (a) surface temperature (K), (b) 850hPa winds ( $\text{ms}^{-1}$ ) and (c) precipitation ( $\text{mm}\cdot\text{day}^{-1}$ ). Unit wind vector is  $5\text{ms}^{-1}$ . Speckling on the precipitation difference (c) indicates significance at the 95% level using a student t-test.

lies in the precipitation response, where the standard model (Fig. 1f) shows a wetter ITCZ and SPCZ, whereas HadCM3IPFA (Fig. 2f) indicates increased precipitation in a band from the SPCZ in the west across to the central ITCZ, and further increases in a band in the central Pacific north of the equator. This may be related to the spatial pattern of applied flux adjustments (TIS05 Fig. 2), which preferentially increases SST in the central equatorial Pacific. Against a background of the warmer climate, this provides warmer mean SSTs and hence increased precipitation due to the non-linear nature of the Clausius-Clapeyron relationship.

In Fig. 3, the effects of flux adjustments alone at  $2 \times \text{CO}_2$  are considered and can be compared to those at  $1 \times \text{CO}_2$  (see TIS05, Fig. 3). The patterns of mean surface temperature change forced by equatorial flux adjustments are qualitatively similar at  $1 \times \text{CO}_2$  and  $2 \times \text{CO}_2$ , reflecting the largely linear response of the ocean surface to identical heat flux anomalies. The wind response is also similar, although the reduction in the monsoon jet over the Arabian Sea is less obvious. This is likely because the temperature changes induced by the anomalous fluxes are taking place against a warmer background state than at  $1 \times \text{CO}_2$ , so their influence on the absolute meridional temperature gradient across the Arabian Sea and Indian subcontinent will be less marked.

In summary, it is possible that the systematic bias in tropical ocean temperatures (which is predominantly cold in the central Pacific and too warm around the Maritime Continent) cause HadCM3 to underestimate the effects of increased greenhouse gas forcing. These biases in the basic state may also propagate through to the interannual variability, as described in TIS05, and thus lead to greater uncertainty in the predictability of monsoon systems in the future climate.

### (c) *Quantifying precipitation changes over India*

Now that both future-climate integrations have been introduced, we can make a quantitative estimate of the precipitation increase over the Indian region. First we consider the Indian land

TABLE 1. DAILY RAINFALL TOTALS ( $\text{mm}\cdot\text{day}^{-1}$ ) DURING THE JJAS SEASON OVER THE INDIAN LAND SURFACE, AS DEFINED BY GADGIL AND SAJANI (1998).

CO <sub>2</sub>	CMAP	HadCM3	HadCM3IPFA	$\Delta\text{rain}_{\text{FA}}$
observed	4.60	-	-	-
$1 \times \text{CO}_2$	-	4.29	3.41	-20.5%
$2 \times \text{CO}_2$	-	4.44	3.84	-13.5%
$\Delta\text{rain}_{\text{CO}_2}$	-	+3.5%	+12.7%	-

TABLE 2. DAILY RAINFALL TOTALS ( $\text{mm}\cdot\text{day}^{-1}$ ) DURING THE JJAS SEASON OVER THE  $5 - 40^\circ\text{N}$ ,  $60 - 100^\circ\text{E}$  REGION AFTER MEEHL AND ARBLASTER (2003).

CO <sub>2</sub>	CMAP	HadCM3	HadCM3IPFA	$\Delta\text{rain}_{\text{FA}}$
observed	4.79	-	-	-
$1 \times \text{CO}_2$	-	5.17	4.76	-7.9%
$2 \times \text{CO}_2$	-	5.44	5.17	-4.9%
$\Delta\text{rain}_{\text{CO}_2}$	-	+5.2%	+8.5%	-

surface gridpoints, as defined by Gadgil and Sajani (1998)<sup>‡</sup>. The summer monsoon season (JJAS) daily rainfall totals are shown in Table 1, together with differences to the  $1 \times \text{CO}_2$  scenarios. The observed values are taken from 1979–2004 CMAP data, bilinearly interpolated to the model grid prior to calculation. Illustrating the difficulty in modelling monsoon precipitation, HadCM3 is seen to overestimate monsoon rainfall over the broad region despite underestimating it over the Indian land surface. Increased greenhouse gas forcing causes an increase in monsoon season precipitation, between +3.5 and +12.7% depending on model version, the flux adjustments having a very significant impact. This reinforces the suggestion that systematic errors in HadCM3 may be masking the true impact of increased CO<sub>2</sub> concentration. It is also interesting to note the effect of flux adjustment alone for a given greenhouse gas concentration. The excessive summer monsoon rainfall in HadCM3 (described in TIS05) is reduced by 20.5% after flux adjustment. Removing the same systematic errors has a smaller but still significant effect in the future climate (−13.5%).

For comparative purposes, and to give an idea of the broader behaviour, seasonal rainfall has also been measured over a larger region covering India, the Bay of Bengal, Arabian Sea and the Himalaya as well as other parts of Southeast Asia. The region is  $5 - 40^\circ\text{N}$ ,  $60 - 100^\circ\text{E}$ , after Meehl and Arblaster (2003), who noted a change in rainfall of +9% during JJA when carbon dioxide concentrations were doubled. The response to increased greenhouse gas concentration in HadCM3, as shown in Table 2, is +5.2 to +8.5% for JJAS (+4.6 to +8.4% for JJA, not shown), and therefore consistent with the results of Meehl and Arblaster (2003).

#### 4. Climate change and monsoon variability

As well as changes in the mean climate, future variability of the monsoon will be of critical importance. Any increase in the spread around the mean will require societies to become increasingly adaptable. Increased monsoon variability in a warmer climate is a common finding of modelling studies, and was first noted by Meehl and Washington (1993) in an early coupled model. In more recent studies at higher resolution, Kitoh *et al.* (1997) and May (2004) found increased monsoon variability in transient and  $2 \times \text{CO}_2$  experiments respectively. As well as All-India Rainfall, the dynamical monsoon index (DMI, Webster and Yang 1992) has been used as a measure of the interannual variability of the monsoon over the broad South Asia region. The index is defined as the anomalous zonal windshear during summer (JJAS),

$$\text{DMI} = \overbrace{(U_{850} - \bar{U}_{850}) - (U_{200} - \bar{U}_{200})} \quad (1)$$

where the overbrace indicates averaging over the  $40 - 110^\circ\text{E}$ ,  $5 - 20^\circ\text{N}$  region. We can compare the two measures of monsoon strength in summer by taking their correlation. Table 3 shows that increasing greenhouse gas concentration results in increased coupling between the modelled monsoon wind

<sup>‡</sup> These are 27 land-surface gridpoints within  $10 - 27.5^\circ\text{N}$ ,  $71.25 - 86.25^\circ\text{E}$ . Coordinates refer to gridpoint centers.

TABLE 3. CORRELATION BETWEEN DMI AND AIR

CO <sub>2</sub>	HadCM3	HadCM3IPFA	ERA-40/AIR
observed	-	-	0.55
1 × CO <sub>2</sub>	0.25	0.47	-
2 × CO <sub>2</sub>	0.40	0.58	-

Correlation between the summer (JJAS) dynamical monsoon index (DMI) and the All-India Rainfall (AIR) for the two models under 1 × CO<sub>2</sub> and 2 × CO<sub>2</sub> conditions, and the reanalysis/observations.




Figure 4. The summer (JJAS) dynamical monsoon index as described in Eq. (1) for (a) HadCM3 2 × CO<sub>2</sub> and (b) HadCM3IPFA 2 × CO<sub>2</sub>. Note that the years in the two model runs are entirely independent.

strength and precipitation. This suggests that in future scenarios the DMI is increasingly useful as an indicator of monsoon strength, although the reason for this is unclear.

The DMI is shown in Fig. 4 for both versions of the model under 2 × CO<sub>2</sub> conditions. A striking feature is the pronounced biennial monsoon behaviour in the HadCM3IPFA 2 × CO<sub>2</sub> integration, *i.e.*, a weak monsoon being followed by a strong monsoon and vice-versa. This tendency of the monsoon in HadCM3IPFA 2 × CO<sub>2</sub> to exhibit tropospheric biennial oscillation (TBO) behaviour (as described in Meehl (1997) and Ogasawara *et al.* (1999) amongst others) will be studied in more detail in part B of this paper (Turner *et al.* 2007). Both models show strong extremes, and probability density functions of the data (not shown) indicate reduced occurrence of conditions close to the mean when compared to integrations at 1 × CO<sub>2</sub>. This result is independent of the index used to measure monsoon strength.

Table 4 summarizes the interannual standard deviation of the DMI, which reaches 1.51ms<sup>-1</sup> in HadCM3 2 × CO<sub>2</sub>. This represents an increase of nearly a quarter over the equivalent 1 × CO<sub>2</sub> integration, but is still lower than that for the ERA-40 reanalysis. Using flux adjustments variability is also increased, as in the 1 × CO<sub>2</sub> scenarios of TIS05, but the response to 2 × CO<sub>2</sub> is not so marked. Consideration of precipitation variability measured over the Indian land surface (defined earlier) yields consistent increases for doubled CO<sub>2</sub>, but apparently little change when flux adjustments are applied. When considered as a percentage of the seasonal total rainfall however, (120days × daily rate values in Table 1) precipitation variability is still increased with flux adjustments.

The effect of both flux adjustments and CO<sub>2</sub> doubling on monsoon variability is one of more frequent and stronger extremes. Further evidence for this can be seen in the following composite analysis, using the DMI to delineate strong and weak monsoons over the broad South Asia region. A composite difference of strong (DMI > 0.5σ) and weak (< -0.5σ) summer monsoon seasons (JJAS) is shown for precipitation and 850hPa winds in Fig. 5, and for surface temperature in Fig. 6. The wind difference pattern for HadCM3 2 × CO<sub>2</sub> is similar to that at 1 × CO<sub>2</sub> (see TIS05 Fig. 7), although the cyclonic anomaly over the Bay of Bengal, the monsoon flow, and the anomalous inflow into the Maritime Continent are all stronger. Associated with these circulation patterns are precipitation

TABLE 4. INTERANNUAL STANDARD DEVIATION OF SUMMER MONSOON STATISTICS

CO <sub>2</sub>	HadCM3	HadCM3IPFA	ERA-40/CMAP
observed	-	-	1.60 (84.3)
1 × CO <sub>2</sub>	1.22 (111.0)	2.05 (109.8)	-
2 × CO <sub>2</sub>	1.51 (126.0)	2.17 (129.1)	-

Interannual standard deviation ( $\sigma$ ) of the summer DMI (AIR) in HadCM3 and HadCM3IPFA under 1 × CO<sub>2</sub> and 2 × CO<sub>2</sub> conditions. Units are ms<sup>-1</sup> (mm). Comparisons are with ERA-40 (CMAP) over the 1958-1997 (1979-2004) periods.




Figure 5. Strong minus weak year composite difference plots of summer (JJAS) lower tropospheric (850hPa) winds and precipitation for (a) HadCM3 2 × CO<sub>2</sub> and (b) HadCM3IPFA 2 × CO<sub>2</sub>. Unit wind vectors are 3ms<sup>-1</sup>.

increases, dynamically strong monsoons now being wetter than weak monsoons by 2mm.day<sup>-1</sup> in the west of India, and up to 3mm.day<sup>-1</sup> in the Bay of Bengal, whilst the Maritime Continent and the Indian Ocean to the west of Sumatra receive a further 4mm.day<sup>-1</sup>. The drier region associated with the anomalously strong Pacific trades is down by 6mm.day<sup>-1</sup> during strong monsoon years. The West African monsoon winds are noticeably stronger in HadCM3 2 × CO<sub>2</sub> during strong Asian summer monsoon years. For HadCM3IPFA 2 × CO<sub>2</sub> (Fig. 5b), as at 1 × CO<sub>2</sub>, the flux adjustments have caused an increase in variability in the Indo-Pacific basins, although the same systematic patterns are evident as in the standard version of HadCM3. Strong Asian summer monsoons are now even wetter over the Arabian Sea, India, and the Bay of Bengal. The anomalous cyclonic flow just south west of Sumatra means that strong monsoon summers are 6mm.day<sup>-1</sup> wetter there than weak summers. There is now evidence of additional precipitation in the Gulf of Guinea, part of the West African monsoon, coinciding with the anomalous westerlies across the continent.

The monsoon difference composite for surface temperature (Fig. 6) shows increased variability in the central equatorial Pacific when compared to 1 × CO<sub>2</sub> (TIS05 Fig. 8): a 1.5°C range in HadCM3 2 × CO<sub>2</sub> and 4°C in HadCM3IPFA 2 × CO<sub>2</sub>. This suggests increased coupling between the monsoon and ENSO at 2 × CO<sub>2</sub>. In the Indian Ocean, flux adjustments cause strong monsoons




Figure 6. Strong minus weak year composite difference plots of summer (JJAS) surface temperature (K) for (a) HadCM3  $2 \times \text{CO}_2$  and (b) HadCM3IPFA  $2 \times \text{CO}_2$ . Negative differences are shaded whilst dashed contours show positive differences. Contour interval is 0.5K.

to be increasingly associated with warm SSTs west of Sumatra, the difference rising from  $1.2^\circ\text{C}$  in HadCM3 to  $1.7^\circ\text{C}$  in HadCM3IPFA. This enhanced coupling between the monsoon and Indian Ocean will be examined in more detail in part B.

## 5. ENSO variability and climate change

Before considering monsoon predictability, interannual variability in the tropical Pacific, to which variations in the monsoon are tied (*e.g.*, Torrence and Webster 1999), will be discussed. Guilyardi (2006) studied the IPCC AR4 model integrations<sup>§</sup> and noted that those models which featured the most realistic mixture of mechanisms when compared to observations displayed an increase in ENSO amplitude in future greenhouse warming scenarios. By considering how models represent different feedback loops in the generation of El Niño-La Niña anomalies, van Oldenborgh *et al.* (2006) demonstrate slight increases, decreases and null responses in ENSO variation, although none lie outside the range of observed decadal variability. Merryfield (2006) also found no consensus on ENSO amplitude change in the IPCC AR4 integrations.

One suggested mechanism for ENSO amplitude change lies in the vertical intensity of the thermocline. Timmermann *et al.* (1999) found much stronger ENSO variability when the ECHAM4/OPYC3 coupled model was subject to IS92a forcing, as did Collins (2000a) in HadCM2, attributable to the increased vertical temperature gradient. This makes temperatures on the ocean surface more prone to variation via thermocline movements. Using similar reasoning, Meehl *et al.* (2006) found that under strong greenhouse forcing, the PCM and CCSM3 coupled models both featured significantly decreased ENSO amplitude, attributable to the warmer but more diffuse thermocline, in turn related to reduced wind driven meridional overturning cells in the upper ocean. Thus the response of ENSO variability to increased greenhouse gas forcing is model dependent.

As shown in Table 5, El Niño amplitude measured as the standard deviation of the Niño-3 index increases under  $2 \times \text{CO}_2$  in both model versions. This is consistent with the increased vertical intensification of the thermocline described in section 3. Increased ENSO variability, when coupled through the atmospheric teleconnection, would provide for greater monsoon variability as witnessed in the previous section. However, other studies have shown the basic state in the Pacific itself to

<sup>§</sup> The WCRP CMIP3 multi-model dataset archived at PCMDI.


TABLE 5. SST ANOMALY AMPLITUDE ( $\sigma$ ) OF THE NIÑO-3 REGION.

CO <sub>2</sub>	HadCM3	HadCM3IPFA	ERA-40
observed	-	-	0.85
1 × CO <sub>2</sub>	0.94	1.21	-
2 × CO <sub>2</sub>	1.05	1.32	-

Units are °C

TABLE 6. CORRELATIONS BETWEEN MONSOON STRENGTH AND MEAN PACIFIC SST OR VARIABILITY IN 11-YEAR WINDOW.

correlation	HadCM3 1 × CO <sub>2</sub>	HadCM3IPFA 1 × CO <sub>2</sub>	HadCM3 2 × CO <sub>2</sub>	HadCM3IPFA 2 × CO <sub>2</sub>
$\sigma(\text{DMI})$ vs. $\text{mean}(\text{Niño-3})$	-0.13	-0.13	-0.08	-0.09
$\sigma(\text{DMI})$ vs. $\sigma(\text{Niño-3})$	0.80	0.88	0.86	0.93

Above  $\sim 0.6$  correlations are significant at the 95% level.Figure 7. Normalized timeseries of Niño-3 SST anomalies at  $2 \times \text{CO}_2$  for (a) HadCM3 and (b) HadCM3IPFA.

influence monsoon variability. Meehl and Arblaster (2003) simulated future greenhouse warming scenarios by imposing fixed warm SST anomalies across the Pacific basin upon a dataset of observed SSTs forcing an atmosphere-only GCM. Stronger interannual variation of the monsoon was noted in their experiment, the warmer basic state causing greater variations in latent heat flux variability to be transmitted via the Walker circulation to the monsoon. In this model framework, this hypothesis is tested by taking correlations between monsoon variability (the standard deviation of the DMI), Pacific variability (Niño-3 SST anomaly standard deviation) and Pacific mean temperatures (Niño-3 mean SST). Each is measured through an 11-year moving window. These correlations are shown for the Niño-3 region in Table 6, which indicates that on decadal timescales at least, correlations between monsoon variability and mean SST in the Pacific are insignificant when compared to correlations with Pacific SST variability. The same result is found if the Niño-3.4 or Niño-4 regions are used instead.

Figure 7 shows the timeseries of Niño-3 SST anomalies in both models at  $2 \times \text{CO}_2$ . In HadCM3, behaviour seems reasonable, encompassing interspersed small and large El Niño-La Niña events separated by a range of periods. This integration shares similar characteristics to  $1 \times \text{CO}_2$  data and observations (not shown). In HadCM3IPFA, however, distinct regimes are present, encompassing a noticeable biennial tendency also evident in the monsoon (Fig. 4). It is of interest that the biennial tendency of the monsoon-ENSO system in HadCM3IPFA is also present in some other IPCC AR4




Figure 8. Correlation between Niño-3 SST and summer (JJAS) (a) DMI or (b) Indian rainfall plotted against lag time. In (b), ERA40 vs AIR refers to the All-India Rainfall precipitation data being correlated with ERA-40 SSTs. Correlations are significant in all but 5% of cases when  $|r| > 0.31$  for observations,  $|r| > 0.2$  for model integrations. Thin (thick) lines are  $1 \times \text{CO}_2$  ( $2 \times \text{CO}_2$ ).

models (Nanjundiah *et al.* 2005). These regimes and the tendency of the model ENSO to oscillate at a higher frequency will be discussed in part B of this paper.

## 6. The changing monsoon-ENSO teleconnection

### (a) Mean relationship

Despite the predicted increase in mean monsoon precipitation, the projected increased interannual variability will place new demands on managing socio-economic impacts. Thus it is now pertinent to consider the monsoon-ENSO teleconnection in these future climate scenarios, and whether predictability on seasonal to interannual timescales will change. Figure 8 shows the lag-correlations between Niño-3 SST anomalies and the DMI and Indian Rainfall. Looking first at monsoon dynamics (Fig. 8a), the teleconnection is strengthened under climate change in both versions of the model, suggesting that the Asian summer monsoon may become more predictable under  $2 \times \text{CO}_2$  conditions. In HadCM3, the timing of the Niño-3 teleconnection maximum is later and comparable to present day reanalysis. Looking at the rainfall relation in Fig. 8(b), however, the impact of climate change is less obvious. The HadCM3 teleconnection seems to weaken, whereas in HadCM3IPFA it stays at broadly the same strength during the summer and autumn. Ashrit *et al.* (2003) found no systematic change to the monsoon-ENSO teleconnection in their transient climate change study, indeed they noted the Niño-3 relation with the DMI to be quite robust, consistent with our study. Other indices however, such as meridional wind shear over the Indian sector, were less stable. In their study of the observed east Pacific SST and Indian rainfall records, Torrence and Webster (1999) found high coherence between the two regions during periods of high ENSO and high monsoon variability. This is confirmed by our strengthened dynamical teleconnections during  $2 \times \text{CO}_2$  forcing, concurrent with enhanced variability in both sectors. Indeed Hu *et al.* (2000) also noticed increased ENSO variability in a transient simulation of a future climate, connected to higher monsoon variability.

One can also verify the effect of flux adjustments alone on the teleconnection. Figure 8 demonstrates that systematic biases in the equatorial oceans have a much more significant impact on the strength of the monsoon-ENSO teleconnection than increased greenhouse forcing. Considering the composite evolution of equatorial Pacific SSTs during El Niño events from each of the future climate scenarios (not shown), the same argument holds as in the  $1 \times \text{CO}_2$  integrations. Flux adjustments shift the warmest waters further east, repositioning the ascending branch of the Pacific Walker circulation such that ENSO anomalies are more easily communicated from the Niño-3 region, strengthening the teleconnection in HadCM3IPFA (see TIS05).

The teleconnections of the HadCM3IPFA  $2 \times \text{CO}_2$  integration in Fig. 8 exhibit an interesting property prior to monsoon development, with strong positive correlations at lead times of 12 to 5 months, reaching  $+0.4$  for the DMI and  $+0.25$  for Indian rainfall. These suggest that east Pacific warmings during one summer are followed by strong Asian monsoons the next, consistent with correlations in the observed data. The strong biennial component to monsoon and ENSO variability (Figs. 4 and 7 respectively) is thus evident in the teleconnection, although when periods of biennial




Figure 9. 21-year moving instantaneous correlation between summer (JJAS) Niño-3 region SSTs and (a) DMI or (b) Indian rainfall in HadCM3 and HadCM3IPFA integrations at  $1 \times \text{CO}_2$  and  $2 \times \text{CO}_2$ . Correlations are significant in all but 5% of cases when  $r < -0.43$ . Thin (thick) lines are for  $1 \times \text{CO}_2$  ( $2 \times \text{CO}_2$ ) integrations.

ENSO variation are removed from the data, the dynamical teleconnection is still stronger during JJAS than in HadCM3IPFA  $1 \times \text{CO}_2$  whereas the rainfall teleconnection is weakened. Variations in the monsoon-ENSO system under different ENSO regimes will be shown in part B of this paper.

#### (b) Interdecadal relationship

Recent weakening of the monsoon-ENSO relationship in the observed record raises interesting questions for monsoon prediction. Indeed the 1997–98 El Niño was the largest on record, yet a normal monsoon prevailed over India, the country receiving 102% of the seasonal average rainfall (Slingo and Annamalai 2000). Whilst there is some evidence to suggest the recent weakening is associated with anthropogenic influences (*e.g.*, Kumar *et al.* 1999) this is by no means certain and does not explain earlier changes in the record.

The change in teleconnection strength may also relate to changes in observed ENSO modal behaviour occurring over the same period. Indeed Slingo and Annamalai (2000) suggest that very strong El Niños such as 1997 may lead to different wind regimes acting on the monsoon system, modulating the local Hadley circulation in such a way as to favour strong monsoon conditions. Some recent studies (*e.g.*, Annamalai and Liu 2005) note that El Niño events during different periods (before and after 1976) have quite a different impact on the monsoon. Others, such as Annamalai *et al.* (2006), which includes more recent observations, suggest that the relationship between El Niño and the monsoon may be strengthening again. The authors cite the deficient monsoons of 2002 (–19%) and 2004 (–13%) during moderate Pacific warmings.

To explore interdecadal changes in ENSO-monsoon teleconnections, moving correlations for summer monsoon statistics versus Niño-3 timeseries are shown for all model integrations in Fig. 9. Most noticeable is the large variation in teleconnection strength through each integration, despite fixed  $\text{CO}_2$  concentrations. Correlations are rarely significant using either monsoon measure in the HadCM3  $1 \times \text{CO}_2$  integration due to poor representation of the basic state (TIS05). In the flux adjusted models (at  $1 \times \text{CO}_2$  and  $2 \times \text{CO}_2$ ) there is much greater correspondence between the strengths of dynamic and rainfall correlations as noted in Table 4.

The stability of teleconnections over the entirety of model integrations has also been studied by Annamalai *et al.* (2006). These authors used 20th century  $\text{CO}_2$  forcing simulations of the IPCC AR4 and found the GFDL- CM- 2.0, GFDL- CM- 2.1 and MRI models to have a range of correlation variation similar to that observed. The moving teleconnections did not match the phase changes seen in observations however, with no noticeable weakening trend. This suggests that climate change may not be the cause of recent weakening in the teleconnection (as suggested by some authors *e.g.*, Kumar *et al.* 1999; Ashrit *et al.* 2005). Indeed whilst recent observed monsoon-ENSO teleconnections seem to have declined, the modelled climate change response has been one of a generally robust relationship (in this and other studies). Ashrit *et al.* (2003) also suggest that recent weakening is more likely related to internal variability of the monsoon-ENSO system. Our study therefore confirms the view that internal variability has a more significant impact on changing monsoon-ENSO teleconnections than climate change.

## 7. Conclusions

The doubling of carbon dioxide concentration in HadCM3 has had several impacts on the mean climate of the coupled Indo-Pacific system. A slight northward shift in the Somali Jet, accompanied with significant increases in seasonal precipitation in the Asian summer monsoon region, amounts to an increase in mean monsoon strength. The increased monsoon precipitation is due to increased moisture convergence over the warmer Indian Ocean, and occurs despite the effects of El Niño-like warming in the tropical Pacific Ocean. There are also significant and consistent increases in precipitation in the Pacific ITCZ from the Philippines to 120°W. The system of limited-area flux adjustments affect the mean climate in the same way as under  $1 \times \text{CO}_2$  conditions, although the response to greenhouse gas increase in HadCM3IPFA is stronger in all mean fields, with an increase of nearly 13% in monsoon precipitation. This illustrates the fundamental importance of reproducing the correct basic state conditions in the coupled system.

The subsurface response to  $\text{CO}_2$  doubling bears a signature of the El Niño-like surface change, resulting in an increased vertical gradient of the thermocline. This contributes to increased SST variability in the east Pacific, a fundamental component of ENSO. Interannual variability of the monsoon increases at  $2 \times \text{CO}_2$ , and rises further with flux adjustments, when the monsoon also exhibits strong biennial behaviour.

Monsoon dynamics are increasingly predictable when using ENSO indicators in the future climate, and the monsoon precipitation teleconnection to the east Pacific remains generally robust. A consequence of this is the increased correspondence between monsoon dynamics and precipitation at  $2 \times \text{CO}_2$ , although the complex relationship between monsoon precipitation and dynamics in GCMs deserves further scrutiny. The flux adjusted model sees a stronger monsoon-ENSO teleconnection using both measures, as under  $1 \times \text{CO}_2$ . This demonstrates the significant impact of model bias on monsoon predictability, an impact greater than climate change itself. The monsoon-ENSO teleconnection is found however to undergo variations on interdecadal timescales, despite the absence of transient external forcing. These variations are of similar amplitude to those in observations and thus changes to the mean teleconnection due to systematic errors or increased greenhouse forcing may lie within this range. Hence there is still much to understand about interdecadal and climatic changes in the coupled monsoon-ENSO system, which may require much longer GCM integrations to be performed.

The pre-season biennial character to the monsoon-ENSO teleconnections in HadCM3IPFA, which indicates the crucial link between the biennial monsoon variability and biennial ENSO in HadCM3IPFA at  $2 \times \text{CO}_2$ , will be further explored in part B of this paper.

## Acknowledgements

A. G. Turner was supported through a PhD studentship provided by NERC's National Centre for Atmospheric Science (NCAS), and is now funded via the EU-ENSEMBLES project. Julia Slingo and Pete Inness are members of NCAS. The authors wish to thank the Associate Editor Richard Seager, Jerry Meehl, and a further anonymous reviewer who have provided useful comments which helped improve this manuscript. In addition we are grateful to Jason Lowe of the Hadley Centre Atmosphere for providing atmosphere and ocean start dumps; the British Atmospheric Data Centre for supplying the ERA-40 dataset; The Climate Prediction Center for CMAP data; the Indian Institute of Tropical Meteorology, Pune, for their All-India Rainfall. Computing resources were provided by CSAR for running the Unified Model.

## References

- Annamalai, H., Hamilton, K. and Sperber, K. R. (2006) 'South Asian Summer Monsoon and its Relationship with ENSO in the IPCC AR4 simulation'. *submitted to J. Climate*
- Annamalai, H. and Liu, P. (2005) Response of the Asian summer monsoon to changes in El Niño properties. *Q. J. R. Meteorol. Soc.*, **131**, 805–831
- Ashrit, R., Douville, H. and Kumar, K. (2003) Response of the Indian Monsoon and ENSO-monsoon teleconnections to enhanced greenhouse effect in the CNRM coupled model. *J. Meteorol. Soc. Jpn.*, **81**, 779–803
- Ashrit, R. G., Kitoh, A. and Yukimoto, S. (2005) Transient Response of ENSO-Monsoon Teleconnection in MRI-CGCM2.2 Climate Change Simulations. *J. Meteorol. Soc. Jpn.*, **83**, 273–291

- Collins, M. (2000a) The El Niño-Southern Oscillation in the Second Hadley Centre Coupled Model and Its response to Greenhouse warming. *J. Climate*, **13**, 1299–1312
- Collins, M. (2000b) Understanding uncertainties in the response of ENSO to greenhouse warming. *Geophys. Res. Lett.*, **27**, 3509–3512
- Collins, M. and The CMIP Modelling Groups (2005) El Niño- or La Niña-like climate change?. *Clim. Dyn.*, **25**, 89–104
- Federov, A. V. and Philander, S. G. (2000) Is El Niño Changing?. *Science*, **288**, 1997–2002
- Gadgil, S. and Sajani, S. (1998) Monsoon precipitation in the AMIP runs. *Clim. Dyn.*, **14**, 659–689
- Gordon, C., Cooper, C., Senior, C. A., Banks, H., Gregory, J. M., Johns, T. C., Mitchell, J. F. B. and Wood, R. A. (2000) The simulation of SST, sea ice extents and ocean heat transports in a version of the Hadley Centre coupled model without flux adjustments. *Clim. Dyn.*, **16**, 147–168
- Guilyardi, E. (2006) El Niño- mean state - seasonal cycle interactions in a multi-model ensemble. *Clim. Dyn.*, **26**, 329–348
- Hu, Z.-Z., Latif, M., Roeckner, E. and Bengtsson, L. (2000) Intensified Asian summer monsoon and its variability in a coupled model forced by increasing greenhouse gas concentrations. *Geophys. Res. Lett.*, **27**, 2618–2684
- Intergovernmental Panel on Climate Change (2001) *Climate Change 2001: The scientific basis*. Contribution of Working Group I to the Third Assessment Report, Eds. J. T. Houghton, Y. Ding, D. J. Griggs, M. Noguer, P.J. van der Linden, X. Dai, K. Maskell and C. A. Johnson, Cambridge University Press, UK
- Inness, P. M., Slingo, J. M., Woolnough, S., Neale, R., and Pope, V. (2001) Organization of tropical convection in a GCM with varying vertical resolution: Implications for the simulation of the Madden-Julian Oscillation. *Clim. Dyn.*, **17**, 777–793
- Inness, P. M., Slingo, J. M., Guilyardi, E. and Cole, J. (2003) Simulation of the Madden-Julian Oscillation in a coupled general circulation model. Part II: The role of the basic state. *J. Climate*, **16**, 365–382
- Johns, T. C., Gregory, J. M., Ingram, W. J., Johnson, C. E., Jones, A., Lowe, J. A., Mitchell, J. F. B., Roberts, D. L., Sexton, D. M. H., Stevenson, D. S., Tett, S. F. B. and Woodage, M. J. (2003) Anthropogenic climate change for 1860 to 2100 simulated with the HadCM3 model under updated emissions scenarios. *Clim. Dyn.*, **20**, 583–612
- Kitoh, A., Yukimoto, S., Noda, A. and Motoi, T. (1997) Simulated Changes in the Asian Summer Monsoon at Times of Increased Atmospheric CO<sub>2</sub>. *J. Meteorol. Soc. Jpn.*, **75**, 1019–1031
- Krishna Kumar, K., Rajagopalan, B. and Cane, M. A. (1999) On the Weakening Relationship Between the Indian Monsoon and ENSO. *Science*, **284**, 2156–2159
- Latif, M., Sperber, K., Arblaster, J., Braconnot, P., Chen, D., Colman, A., Cubasch, U., Cooper, C., Delecluse, P., DeWitt, D., Fairhead, L., Flato, G., Hogan, T., Ji, M., Kimoto, M., Kitoh, A., Knutson, T., Le Treut, H., Li, T., Manabe, S., Marti, O., Mechoso, C., Meehl, G., Power, S., Roeckner, E., Sirven, J., Terray, L., Vintzileos, A., Voss, R., Wang, B., Washington, W., Yoshikawa, I., Yu, J. and Zebiak, S. (2001) ENSIP: the El Niño simulation intercomparison project. *Clim. Dyn.*, **18**, 255–276
- Martin, G. M., Arpe, K., Chauvin, F., Ferranti, L., Maynard, K., Polcher, J., Stephenson, D. B. and Tschuck, P. (2000) Simulation of the Asian Summer Monsoon in Five European General Circulation Models. *Atmos. Sci. Lett.*, **1**, 37–55
- May, W. (2004) Potential future changes in the Indian summer monsoon due to greenhouse warming: analysis of mechanisms in a global time-slice experiment. *Clim. Dyn.*, **22**, 389–414
- Meehl, G. A. (1997) The South Asian Monsoon and the Tropospheric Biennial Oscillation. *J. Climate*, **10**, 1921–1943
- Meehl, G. A., Collins, W. D., Boville, B. A., Kiehl, J. T., Wigley, T. M. L. and Arblaster, J. M. (2000) Response of the NCAR climate system model to increased CO<sub>2</sub> and the role of physical processes. *J. Climate*, **13**, 1879–1898
- Meehl, G. A. and Arblaster, J. M. (2003) Mechanisms for projected future changes in south Asian monsoon precipitation. *Clim. Dyn.*, **21**, 659–675
- Meehl, G. A. and Tang, H. and Branstator, G. (2006) Future changes of El Niño in two coupled climate models. *Clim. Dyn.*, **26**, 549–566
- Meehl, G. A. and Washington, W. M. (1993) South Asian Summer Monsoon Variability in a Model with Doubled Atmospheric Carbon Dioxide Concentration. *Science*, **260**, 1101–1104
- Meehl, G. A. and Washington, W. M. (1996) El Niño-like climate change in a model with increased atmospheric CO<sub>2</sub> concentrations. *Nature*, **382**, 56–60

- Merryfield, W. J. (2006) Changes to ENSO under CO<sub>2</sub> Doubling in a Multimodel Ensemble. *J. Climate*, **19**, 4009–4027
- Nanjundiah, R. S., Vidyunmala, V. and Srinivasan, J. (2005) The impact if increase in CO<sub>2</sub> on the simulation of tropical biennial oscillations (TBO) in 12 coupled general circulation models. *Atmospheric Science Letters*, **6**, 183–191
- Ogasawara, N., Kitoh, A., Yasunari, T. and Noda, A. (1999) Tropospheric Biennial Oscillation of ENSO-Monsoon System in the MRI Coupled GCM. *J. Meteorol. Soc. Japan*, **77**, 1247–1270
- van Oldenborgh, G. J., Philip, S. and Collins, M. (2006) El Niño in a changing climate: A multi-model study. *Ocean Sci. Discuss.*, **2**, 267–298
- Parthasarathy, B., Munot, A. A. and Kothawale, D. R. (1994) All-India Monthly and Seasonal Rainfall Series: 1871-1993. *Theor. Appl. Climatol.*, **49**, 217–224
- Pope, V. D., Gallani, M. L., Rowntree, P. R. and Stratton, R. A. (2000) The impact of new physical parametrizations in the Hadley Centre climate model: HadAM3. *Clim. Dyn.*, **16**, 123-146
- Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., Rowell, D. P., Kent, E. C. and Kaplan, A. (2003) Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century. *J. Geophys. Res.*, **108(D14)**, 4407, doi:10.1029/2002JD002670
- Slingo, J. M. and Annamalai, H. (2000) 1997: The El Niño of the Century and the Response of the Indian Summer Monsoon. *Mon. Weather Rev.*, **128**, 1778–1797
- Spencer, H. and Slingo, J. M. (2003) The simulation of peak and delayed ENSO teleconnections. *J. Climate*, **16**, 1757–1774
- Sperber, K. R. and Palmer, T. N. (1996) Interannual tropical rainfall variability in general circulation model simulations associated with the atmospheric model intercomparison project. *J. Climate*, **9**, 2727–2750
- Sutton, R. T., Dong, D. and Gregory, J. M. (2007) Land/sea warming ratio in response to climate change: IPCC AR4 model results and comparison with observations. *Geophys. Res. Lett.*, **34**, L02701
- Timmermann, A., Oberhuber, J., Bacher, A., Esch, M., Latif, M. and Roeckner, E. (1999) Increased El Niño frequency in a climate model forced by future greenhouse warming. *Nature*, **398**, 694–697
- Timmermann, A., Jin, F.-F. and Collins, M. (2004) Intensification of the annual cycle in the tropical Pacific due to greenhouse warming. *Geophys. Res. Lett.*, **31**, 2727–2750
- Torrence, C. and Webster, P. J. (1999) Interdecadal changes in the ENSO-monsoon system. *J. Climate*, **12**, 2679–2690
- Turner, A. G., Inness, P. M. and Slingo, J. M. (2005) The role of the basic state in the ENSO-monsoon relationship and implications for predictability. *Q. J. R. Meteorol. Soc.*, **131**, 781–804
- Turner, A. G., Inness, P. M. and Slingo, J. M. (2007) ‘The effect of doubled CO<sub>2</sub> and model basic state biases on the monsoon-ENSO system. Part B: Changing ENSO regimes’. *submitted to Q. J. R. Meteorol. Soc.*
- Uppala, S. M., P. W. Kallberg, A. J. Simmons, U. Andrae, V. da Costa Bechtold, M. Fiorino, J. K. Gibson, J. Haseler, A. Hernandez, G.A. Kelly, X. Li, K. Onogi, S. Saarinen, N. Sokka, R. P. Allan, E. Andersson, K. Arpe, M. A. Balmaseda, A. C. M. Beljaars, L. van de Berg, J. Bidlot, N. Bormann, S. Caires, F. Chevallier, A. Dethof, M. Dragosavac, M. Fisher, M. Fuentes, S. Hagemann, E. Holm, B. J. Hoskins, L. Isaksen, P. A. E. M. Janssen, R. Jenne, A. P. McNally, J.-F. Mahfouf, J.-J. Morcrette, N. A. Rayner, R. W. Saunders, P. Simon, A. Sterl, K. E. Trenberth, A. Untch, D. Vasiljevic, P. Viterbo and J. Woollen (2005) The ERA-40 re-analysis. *Q. J. R. Meteorol. Soc.* **131**, 2961–3012
- Webster, P. J. and Yang, S. (1992) Monsoon and ENSO: Selectively interactive systems. *Q. J. R. Meteorol. Soc.*, **118**, 877–926
- Webster, P. J., Magana, V. O., Palmer, T. N., Shukla, J., Tomas, R. A., Yanai, M. and Yasanuri, T. (1998) ‘Monsoons: Processes, predictability, and prospects for prediction’. *J. Geophys. Res.*, **103**, 14451–14510
- Xie, P. and Arkin, P. A. (1997) Global Precipitation: A 17-Year Monthly Analysis Based on gauge Observations, Satellite Estimates, and Numerical Model Outputs. *Bull. Am. Meteorol. Soc.*, **78**, 2539–2559